

Prescription shoes

If you have been supplied with shoes, they will have been made to a prescription. You should follow the instructions your orthotist (the person who designed or prescribed your shoes) or podiatrist gives you. At first you should wear the shoes for short periods around your home, checking for any problems such as redness or a change in skin colour caused by rubbing or pressure. If this occurs, you should contact your podiatrist or orthotist immediately.

When you and the person who prescribed your shoes are happy with the comfort and fit of the first pair, you will be supplied with a second pair. These should be the only shoes you wear. It is important that you wear these shoes inside and outside the house.

Footwear will normally be prescribed with insoles. These are an important part of your prescription, and you should only remove them if your podiatrist or orthotist advises you to. It is important that you check the inside of your shoes every day to make sure no small objects have fallen in, and that there is no damage to the insole or lining.

If you notice any wear or damage to the shoes, you should contact your orthotist or podiatrist so that they can be repaired.

Whoever provided your shoes will advise you about any repairs or alterations to make sure that they still match your prescription.

Socks, stockings and tights

You should always change your socks, stockings or tights every day. They should not have bulky seams. The tops of socks should not be elasticated. Some high-street shops sell suitable products - ask your podiatrist for details. All your socks, stockings and tights should be the right size for your feet.

Remember: if you are not sure about any shoes, socks, stockings or tights you are wearing, or new shoes you have bought, your podiatrist will be happy to advise you.

Individual advice

.....

.....

.....

.....

.....

.....

.....

.....

Your current foot risk level

Your most recent foot screening has shown that your risk of developing foot complications and foot ulcers because of your diabetes is:

Low / Medium / High / In remission

Local contact numbers

Podiatry Department or Foot Protection Team:

.....

Multi-Disciplinary Foot Care Team:

.....

Normal diabetes clinic:

.....

MLT.MODISK.18_08818 16086. Produced by the Scottish Diabetes - Foot Action Group. This leaflet is for all people with a diagnosis of diabetes, including children over the age of 12. Based on the original leaflet produced by the Scottish Diabetes Group - Foot Action Group, with help from service users. Owned by the Royal College of Podiatry © Published date: February 2025. Review date: February 2026. We would welcome your feedback on this leaflet. Please send it to the Royal College of Podiatry at feedback@rcpod.org.uk



Footwear advice to reduce the risk of non-healing foot wounds or amputation

**Diabetes information and
advice to help protect your
life and limbs**

Footwear and diabetes

This leaflet gives you advice about wearing the most appropriate shoes, taking account of the shape of your feet and your diabetes.

Footwear which does not fit properly is one of the most common causes of foot problems and wounds in people with diabetes.

Why is diabetes a risk to your feet?

The development of foot wounds in people with diabetes is serious as they are linked to an increased risk of heart attacks, strokes, amputations of the foot or leg and early death.

Managing your diabetes, cholesterol and blood pressure, stopping smoking, increasing physical activity and managing your weight helps to reduce the risk of these life-and limb-threatening problems.

Note: You may be at further risk of cardiovascular problems if you have a family history of heart disease.

If you develop any of the problems in the list below, it is important that you contact your Multi-disciplinary Foot Care Team, local Podiatry Department or doctor (GP) for advice as soon as possible (within 24 hours). If none of these are available, please contact NHS 111 by ringing 111. They may advise you to go to your local accident and emergency department.

- An accident or injury to a toe or foot
- A change in the colour or shape of your foot, or swelling to a toe or foot
- A change in the temperature of your foot or lower limb
- New or unexplained pain in your foot
- Oozing or odour from a wound on a toe or foot
- A wound or break in the skin to a toe or foot

Any delay in getting advice or treatment can lead to serious problems.

If you discover any breaks in the skin or blisters, cover them with a sterile dressing. Do not burst blisters.

What can I do to reduce my risk of developing problems?

Check your feet every day

Check your feet every time you remove your shoes, stockings or tights and check for any redness or blisters.

Check your shoes

Check the bottom of your shoes before putting them on to make sure that nothing sharp such as a pin, nail or glass has pierced the outer sole. Also, run your hand inside each shoe to check that no small objects such as small stones have fallen in. Any objects that are inside your footwear or have pierced the outer sole can put your feet at risk of injury.

The shape of your feet

Your feet can change shape over time, so when you buy shoes always check they fit properly.

Stopping smoking

If you smoke, you are strongly advised to stop. Smoking affects your circulation, which can increase the likelihood of amputation.

Buying new shoes

Get your feet measured

Get your feet measured and your new shoes fitted properly by a trained professional at a reputable retailer.

Well-fitting shoes

Make sure your shoes are long enough, deep enough and wide enough for your feet. If your shoes are too tight they will press on your toes and can cause pressure-related problems and wounds, which you may not feel if you have reduced feeling in your feet.

When buying new shoes, always try on both shoes. Most people's feet are slightly different sizes.

Buy your own shoes

If possible, you should go to the shoe shop yourself to have the footwear fitted and not ask somebody else to buy your shoes for you.

Wear new shoes around the house

Wear new shoes around the house for short periods (20 minutes) and then check your feet. Look for problems such as redness caused by rubbing or pressure. If this happens, you can usually return shoes to the shop, but only if you have not worn them outside. If there are no signs of redness or rubbing, slowly increase the amount of time you wear the shoes for each day by 20 minutes, building up to a full day.

Advice on new shoes

If you are not sure if your new shoes are suitable, ask your podiatrist for advice before you wear them.

Recommended footwear

Types of shoes

A well-fitting shoe, boot or trainer with laces or a strap fastening will give your feet the best support. These fastenings will help to keep your foot firmly in place inside your footwear and help prevent rubbing.

Heel height

The heel of your footwear should not be more than 2.5 centimetres (1 inch), to reduce excessive pressure on the front of your foot.

Natural materials

Where possible the uppers (top) of your shoes should be made from a natural material such as soft leather. Leather will fit to your foot without causing any rubbing and will help to prevent your feet from swelling. Check the condition of your prescribed footwear and have repairs done when necessary.

Wear and tear

Signs of wear and tear

Check your shoes for signs of wear such as rough edges in seams or linings, and worn soles and heels. Always check your shoes are in good condition and replace them whenever there are signs of wear and tear that can't be repaired.